

ESSENTIAL OIL COMPOSITION AND ANTIOXIDANT, ANTICANCER ACTIVITIES OF ARTEMISIA FRIGIDA WILLD. GROWN IN MONGOLIA

Irekhbayar Jambal^{1*}, Amarjargal Ayurzana², Altantsetseg Shatar³

¹*Department of Chemistry, School of Arts and Sciences, National University of Mongolia (NUM), Baga toiruu, Sukhbaatar District, Ulaanbaatar 14201, Mongolia.*

²*Department of Chemistry, Khovd branch school of National University of Mongolia, Jargalant soum, Khovd aimag, 84140, Mongolia. E-mail: amar731012@yahoo.com*

³*Institute of Chemistry and Chemical Technology, MAS, 4th Building Enkhtaivan Avenue, Ulaanbaatar 210351, Ulaanbaatar, Mongolia.*

***Correspondence Author:**

irekhbayar@num.edu.mn

Abstract:

Medicinal plants have always been considered a healthy source of life for all people. In Mongolia, the use of various medicinal and food plants has a long history. In recent years, interest in plant-derived food additives has grown. This study was designed to determine essential oil chemical composition and evaluate antioxidant, cytotoxic activities of aerial parts serial fractions from *Artemisia frigida* Willd. grown in Mongolia. The chemical composition of essential oil from *A. frigida* Willd. was determined by GC-MS analysis. Among them 40 components (35.07%), 1,8-cineol (27.22%), borneol (3.01%), terpin-4-ol (4.51%), γ -cadinene (7.5%), α -terpineol (2.98%), bornylacetate (1.80%) were found to be the major ones. The ethanol crude extract of *A. frigida* Willd. was suspended in water and consequently fractionated with n-hexane, chloroform, ethyl acetate and butyl alcohol. All the fractions were examined for their antioxidant and cytotoxic activities by using DPPH and MTT assays, respectively. Ethyl acetate fraction showed the highest DPPH radical scavenging activity at a concentration of 50 μ g/ml by 93.17% with the IC₅₀ value of 11.68 μ g/ml. MTT results showed that water fraction demonstrated significant cytotoxicity against A549 - human alveolar basal epithelial cell line, with values of 78.69 and 60.98% (50; 10 μ g/ml; at 48 h). These suggest that the fractions of *A. frigida* Willd. could hold a good potential for use in the pharmaceutical industry.

Keywords: *A. frigida* Willd.; essential oil; antimicrobial; antioxidant, and cytotoxic activities

INTRODUCTION

Several plants were widespread for their many therapeutic and pharmaceutical virtues, especially antioxidant, anti-tumoral, and anti-infectious activities. A big part of the world's population still relies on the benefits of food for the treatment of common illnesses (Zhang, 2004).

Many natural compounds extracted from plants exhibit important biological activities (Jian-Qing Yu, 2011). A number of aromatic medicinal plants used for treating infectious diseases have been mentioned in different phytotherapy manuals due to their availability, fewer side effects, and reduced toxicity. The essential oils of these aromatic plants (Vandendool & Kratz, 1963) are responsible for their fragrance as well as biological properties (Kalemba & Kunicka, 2003). Essential oils are complex mixtures of volatile secondary metabolites that mainly consist of mono- and sesquiterpenes and are responsible for both the fragrant and biological effects of aromatic medicinal plants (Salzer, 1977; Angioni, 2003; Senatore, 2004). An important characteristic of essential oils and their constituents is their hydrophobicity, which enables them to partition in the lipids of bacterial cell membranes and mitochondria, thus disturbing the structures and rendering them more permeable (Sikkema, 1995). Essential oil plants have recently received much attention by the pharmaceutical industry due to their multiple functions, especially their antioxidant, anticancer and antimicrobial activities (Bakkali, 2008; Sylvestre, 2006).

The major constituents of many these oils are phenolic compounds (terpenoids and phenylpropanoids) (Lawrence, 2005). Nevertheless, aromatic plants producing non-phenolic essential oils are also used as spices and in folk remedies as antiseptics. (Daise Lopez-Luts, 2008). Free radicals and reaction oxygen species (ROS) are constantly produced as byproducts in the human body during cell metabolism. These harmful byproducts, if not eliminate immediately, can cause oxidative damage to functional macromolecules such as DNA, proteins, and lipids (Apel, 2004; Tang, 2004).

This increases the chance of occurrence of age-related disorders, cancer, atherosclerosis, neurodegenerative diseases, and inflammation (Tang, 2004; Cai, 2004). Consumption of antioxidants, through diet and supplements, is expected to remove ROS from the living system and provide health benefits. Several studies demonstrated that medicinal plants are a rich source of antioxidant compounds such as phenolics, flavonoids, quinones, vitamins, coumarins, and alkaloids, which can decrease the incidence of oxidative stress and associated diseases (Cai, 2004).

Cancer is a general term applied to malignant diseases characterized by rapid and uncontrolled abnormal cells formation which may mass together to form a growth or proliferate throughout the body and it may progress until it causes death. Medicinal plants are the most exclusive source of life saving drugs for the majority of the world's population. Medicinal herbs have been widely used for treatment of diseases in traditional way for several generations. An interaction between traditional medicine and modern biotechnological tools is to be established towards new drug development. The interference between cell biology, in vitro assays and structural chemistry will be the best way forward to obtain valuable leads. There is considerable scientific evidence to suggest that nutritive and non nutritive plant-based dietary factors can inhibit the process of carcinogenesis effectively.

Cancer chemoprevention involves pharmacologic intervention with synthetic or naturally occurring chemicals to prevent, inhibit or reverse carcinogenesis or prevent the development of invasive cancer. Out of an estimated 250,000 higher plants, less than 1% has been screened pharmacologically (El-Shemy et al., 2007). In recent years, focus on plant research has increased all over the world. Antioxidants are a group of substances that are useful for fighting cancer and other processes that potentially lead to diseases such as atherosclerosis, Alzheimer's, Parkinson's, diabetes and heart disease (Valko et al., 2007). Unlike cytotoxic agents that damage tumor cells, antioxidants act by preventing the onset of cancer during carcinogenesis, and they are generally beneficial to cells. Oxidants such as reactive oxygen species (ROS) that include the superoxide radical, hydroxyl radical, hydroperoxyl radical and nitrogen species (RNS) such as peroxy nitrite and nitric oxide damage macromolecules, including proteins, lipids, enzymes and deoxyribonucleic acid (DNA). To combat these radicals, living organisms produce enzymes (for example, catalase, superoxide dismutase and peroxidase) or rely on nonenzymatic molecules, such as glutathione, cysteine, ascorbic acid, flavonoids and vitamin K for protection (Ahmed, 2012).

Artemisia frigida Willd. belongs to the genus *Artemisia* of *Asteraceae* (*Compositae*). *Artemisia* (*Wormwood*) is a large, diverse genus of plants with between 200-500 species which are mainly found in Asia, Europe and North America (Bora & Sharma, 2011). Among them 105 species grow in Mongolian Forest-Steppe and Desert-Gobi (Grubov, 1982). Many *Artemisia* species, which are known by such common names as mugwort, sagebrush, sagewort, and wormwood, have a vast range of biological activities, including antimalarial, cytotoxic, antifungal, antibacterial, antioxidant, and other useful effects (Bora & Sharma, 2011).

They have a characteristic scent or taste, based on monoterpenes and sesquiterpenes, which in many cases are the reason for their application in Mongolian traditional medicine (Shatar, 1998; Markova, 1985). These herbs are used worldwide in tonic, stomachic and stimulant beverages and as antiphlogistics in antiseptic oils or tinctures applied for the relief of rheumatic pains (Verezovskaya, 1991).

Mongolia is rich in essential oil medicinal plants. Mongolian traditional medicine has long history of more than 2500 years. There are about 60 clans, 200 species, 300 kinds of essential oil plants and 600 kinds of herbal plants have been registered, among of them, 150-200 kinds are commonly used (Shatar, 2000; Ligaa, 2005). Many essential oil plants have not been studied yet. It is notable to investigate their chemical compositions and biological activities by using traditional medicine (Shatar, 1989).

To our knowledge, there are no published reports on the chemical composition, biological activities of the essential oil and various fractions of *A. frigida* Willd. grown in Mongolia. Therefore, the aim of this study was to determine essential oil chemical composition and describe their antioxidant, cytotoxic effects of various extracts from *A. frigida* Willd. grown in Mongolia. Consequently, it is important to develop a better understanding of their mode of biological action for new application in human health.

2. Experimental

2.1. Chemicals

Dimethyl sulfoxide (DMSO), MTT, and DPPH were purchased from Sigma Chemical Company. RPMI 1640 medium, fetal bovine serum, penicillin, and streptomycin were purchased from GIBCO Co. (GIBCO BRL, Grand Island, NY, USA). The human alveolar basal epithelial cell lines (A549) were purchased from Korean cell line bank (KCLB, Korea). All other chemicals were of analytical grade and purchased from Sigma-Aldrich (USA) and DUKSAN Co. (Korea).

2.2. Plant material

Wild plants materials used in this investigation were collected from Eastern-Mongolia, in August 2009. The plants were taxonomically identified by Prof. S.Shatar from the Institute of Chemistry and Chemical Technology (MAS), Mongolia. Voucher specimen has been deposited in the herbarium section of Department of Organic and Food Chemistry, National University of Mongolia (NUM), Ulaanbaatar, Mongolia.

2.3. Extraction of the essential oil

The aerial parts of the freshly collected plants were finely chopped and hydro-distilled for 3 h using a Clevenger-Adams type apparatus (Adams, 1991). The yield of the essential oil produced during the steam distillation was 0.96% (v/w). The oil samples obtained were dehydrated over anhydrous sodium sulphate and stored at 4°C prior to analysis.

2.4. Extraction and fractionation of plant

Different crude extracts of wild and traditional plants as well as, fresh plants were prepared according to Ferrigni et al. (Ferrigni, 1982). The air-dried and powdered whole plant was extracted with 70% ethanol (2 L × 3) using sonicator under room temperature. The resultant extracts were combined and evaporated in a rotary vacuum evaporator (Buchi R-205, Switzerland) at 40°C to afford crude extracts. The ethanol crude extracts were suspended in water and then fractionated successively with n-hexane, chloroform, ethyl acetate and butyl alcohol by using the separation funnel, respectively. All the extracted materials were preserved at -20°C until analysis of biological activities.

2.5. Analysis of essential oil

A Hewlett-Packard HP 6890 gas chromatograph coupled with a DB-5 fused silica capillary column (30 m × 0.25 mm × 0.25 µm film thickness, J&W Scientific) and FID detector was used for the quantitative determination of oil components. Oven temperature was programmed as follows: 50°C for 2 min, rising to 250°C at 5°C/min. Injector temperature: 270°C. Carrier gas: He with a flow rate of 1 ml/min. Split ratio 1:10. The volume injected 1.0 µl of 1% solution (diluted in hexane). Identification of the oil components was based on their retention indices and MS, obtained from GC/MS analysis on a HP 6890/HP5973 equipped with DB-5 fused silica capillary column (30 m × 0.25 mm × 0.25 µm film thickness, J&W Scientific). The EI/MS spectra were recorded at 70eV. The mass range was 35 to 500 m/z. The essential oil components were identified on the basis of their mass data and the comparison of their relative retention index (RRI) (Vandendool & Kratz, 1963) that was obtained using various series of n-alkanes. Their EI-mass spectra were either compared with the NIST/NBS and Wiley library spectra found in the literature (Massada, 1976), or were confirmed by comparison with data published in a reference book (Adams, 2001). The percentages of the individual components were obtained from the relative GC peak areas as shown in Table 1.

2.6. Antioxidant activity measurement

2.6.1. DPPH assay

The free radical scavenging activity (antioxidant activity) has been evaluated using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) method, which is regarded as an easy, reliable, sensitive, and rapid method as compared with other antiradical methods (Moon & Shibamoto, 2009). A review by Moon and Shibamoto also reported that 90% of antioxidant activity measurements are conducted by this method. In vitro antioxidant activities of the plants were evaluated on the basis of their ability to inhibit H₂O₂ induced oxidative stress in yeast cells (*Saccharomyces cerevisiae*). This method has been employed for in vivo antioxidant studies as yeast is the ideal model organism for monitoring the biological mechanism (Azevedo, 2011; Lee, 1999).

The assay was carried out according to the method of Brand-William et al. (Brand-Williams, 1995) with a little modification to investigate the free radical scavenging activity of samples. Briefly, the samples were dissolved in ethanol

at the concentration of 100 mg/ml and then serially diluted by ethanol. On each well of a 96-well plate, 100 µl of samples of different concentration were mixed together with 100 µl of 60 µM DPPH prepared in ethanol. After incubation of 20-30 minutes for reaction, the absorbance of supernatants was measured at 517 nm. Ethanol was used as a negative control and α-tocopherol as a positive control.

The scavenging capacity (SC) of the sample was calculated using the following formula:

$$SC (\%) = [1 - A_s/A_c] \times 100$$

Where, A_s = is the net absorbance of the sample, A_c = is the net absorbance of negative control. The IC_{50} value of a sample is the concentration of sample at which 50% activity of DPPH (absorbance) is inhibited. It was calculated by linear regression.

2.7. Cell lines and culture

The human alveolar basal epithelial cell lines (A549) were purchased from Korean cell line bank. The cells were grown in RPMI 1640 supplemented with 10% fetal bovine serum (FBS), 1% (w/v) glutamine, sodium pyruvate 5%, 100 U/ml penicillin and 100 µg/ml streptomycin. Cells were cultured in a humidified atmosphere at 37°C in 5% CO₂.

2.8. MTT assay

The 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl tetrazoliumbromide (MTT) colorimetric method was used for determining cytotoxic activity of samples as describing by Mosman (Mosman, 1983). The MTT colorimetric assay is an established method of determining viable cell number in proliferation and cytotoxicity studies. This assay is based on the cleavage of the yellow tetrazolium salt, MTT, to form a soluble blue formazan product by mitochondrial enzymes, and the amount of formazan produced is directly proportional to the number of living, not dead cells, present during MTT exposure. Since the MTT assay is rapid, convenient, and economical, it has become a very popular technique for quantification of viable cells in culture (Sylvester, 2011).

In brief, each sample was dissolved in DMSO at concentration of 100 mg/ml and serially diluted into different concentration of 1-100 µg/ml with culture media. On a 96-well plate, 200 µl of cell suspension at density of 2×10^4 (in ml) were seeded. After 24 h incubation at 37°C, 5% CO₂ to allow cell attachment, the culture media were removed and replaced by 200 µl of cultured media containing different concentration of samples and incubated for 24 and 48 h at the same condition. In control wells, the media were added without any samples. Finally, after 4 h of MTT reagent addition (final concentration of 0.5 mg/ml), the formazan crystals formed was resolved in DMSO (200 µl/well) and absorbance were measured at 570 and 630 nm. The cell growth inhibition (GI) of the sample was calculated by the formula:

$$GI (\%) = [1 - A_s/A_c] \times 100$$

Where A_s = was the absorbance in sample wells and A_c = was the absorbance in the control wells.

3. Results and discussion

3.1. Chemical composition of essential oil

The essential oil composition was determined by GC-MS analysis. A total of 40 components were identified, representing 95% of the total oil composition. The terpenoides made up the largest component of the oil and had many representative volatiles. The oxygenated monoterpenes (77.84%), oxygenated sesquiterpenes (10.51%), monoterpene hydrocarbons (6.21%) and whereas sesquiterpenes were weakly represented (5.35%).

The main constituents were found to be camphor (35.07%), 1,8-cineol (27.22%), borneol (3.01%), terpin-4-ol (4.51%), γ-cadinene (7.5%), α-terpineol (2.98%), bornylacetate (1.80%).

These differences might have been derived from local, climatic and seasonal factors. The percentage content of the individual components, the retention times and percent yields are summarized in Table 1.

Table 1: Essential oil chemical components of *A. frigida* Willd. grown in Mongolia

Chemical components	Percentage, %	Chemical components	Percentage, %
α-Thujene	0.07	β-Thujone	0.24
α-Pinene	0.78	(E)-Sabinene hydrate	0.78
Camphene	1.30	Fenchyl acetate	0.09
β-Pinene	0.55	Decanal	0.07
Sabinene	0.21	Camphor	35.07
Limonene	0.22	Linalool	0.25
(Z)-β-Ocymene	0.63	(Z)-Sabinene hydrate	0.18
γ-Terpinene	1.48	Octanol	0.07
p-Cymene	0.63	Isobornyl acetate	0.69
Terpinolene	0.30	Bornyl acetate	1.80
1,8-Cineol	27.22	Terpene-4-ol	4.51

α -Copaene	0.07	α -Terpineol	2.98
β -Caryophyllene	0.08	Borneol	3.01
β -Farnesene	0.12	Geraniol	0.12
Humulene	0.41	(E)-Carveol	0.38
Germacrene-D	1.52	Thymol	0.12
δ -Cadinene	3.15	Caryophyllene oxide	1.00
Hexanol	0.10	(E)-Nerolidol	0.50
1-Octene-3-ol	0.16	α -Terpineol	9.01
Monoterpene hydrocarbons	6.17	Oxygenated monoterpenes	77.84
Sesquiterpene hydrocarbons	5.35	Oxygenated sesquiterpenes	10.51

3.2. Antioxidant activity

DPPH is a stable nitrogen-centered free radical the color of which changes from violet to yellow upon reduction by either the process of hydrogen- or electron-donation. Substances which are able to perform this reaction can be considered as antioxidants and therefore radical scavengers (Brand-Williams, 1995).

Antioxidant activities of the different fractions from *A.frigida* Willd. were tested by the DPPH radical scavenging assay. The effect of antioxidant on DPPH radical scavenging was thought to be due to their hydrogen donating ability or radical scavenging activity. When a solution of DPPH is mixed with that of a substance that can donate a hydrogen atom, then this gives rise to the reduced form DPPH (non radical) with the loss of this violet color (Molyneux, 2004).

Among all fractions the ethyl acetate fraction showed strong free radical scavenging activity in DPPH assay system with IC_{50} value of $11.68 \pm 3.5\%$. The scavenging activity of the various fractions in decreasing order was: Ethyl acetate (EtOAc) > n-hexane (Hex) > n-butanol (BuOH) > ethanol (EtOH) > water (H₂O) > chloroform (CHCl₃) (Fig 1).




Fig 1: Free radical scavenging activity (%) by the DPPH assay of serial fractions of *A.frigida* Willd. at a concentration of 50 µg/ml

3.3. Cytotoxic activity

To investigate the cytotoxic activity of different fractions from *A.frigida* Willd., we evaluated its effect on a selection of A549 (human alveolar basal epithelial) cell lines by MTT assay.

These cell lines were submitted to growing concentrations of *A.frigida* fractions for 24 and 48 hours. As shown in Fig 2, the extracts of plants significantly active against chosen human cancer cell lines tested. The cytotoxic activity of the different fractions in decreasing order was: Ethyl acetate (EtOAc) > water (H₂O) > chloroform (CHCl₃) > ethanol (EtOH) > n-butanol (n-BuOH) (Fig 2).




Fig 2: Cell inhibition (%) of organic fractions from *A.frigida* Willd. against A549 cell line. Cells were treated with the fractions for 48 h at a concentration of 50µg/ml

4. Conclusions

In recent years, interest in plant-derived food additives has grown. Plant extracts might substitute synthetic food antioxidants, which may influence human health when consumed chronically (Martinez-Tome, 2001). The results of our study confirmed the importance of the investigated plants in our nutrition.

This study on essential oil chemical composition and biological activities of *A.frigida* Willd. grown in Mongolia were not well performed before. Essential oils hydrodistilled from *A.frigida* Willd. were found to be rich in camphor, 1,8-cineol, borneol, terpin-4-ol, γ -cadinene, α -terpineol and bornylacetate. The antioxidant activity of the ethyl acetate fraction was moderate than other fractions. The results clearly showed that the n-butanol fraction presented satisfactory cytotoxic activity against A549 human cancer cell lines tested. The results of this work also demonstrate the potential of *A.frigida* Willd. various fractions as a new antioxidant and cytotoxic agents for human health.

Acknowledgements

This work was supported by grants from the KFAS (Korean Foundation for Advanced Studies). We are thankful to Prof. Dr. Burm-Jong Lee, Biohealth Products Research Center, Department of Biomedical Chemistry, Inje University, Republic of Korea, and for his support to carry out this investigation. We are also grateful Prof. Sc.Dr. Shatar Sandui, Institute of Chemistry and Chemical Technology (MAS), highly appreciate for his assistance to collect and help in GC/MS-analysis of essential oil.

References

- [1]. Adams, R.P. (1991). Cedar Wood Oil - Analysis and properties. In: Modern Methods of Plant Analysis: Oils and Waxes. Springer-Verlag Berlin, Germany, 159-173.
- [2]. Azevedo, F.; Marques, F.; Fokt, H.; Oliveira, R.; Johansson, B. (2011). Measuring oxidative DNA damage and DNA repair using the yeast comet assay. *Yeast*, 28, 55-61.
- [3]. Adams, R.P. (2001). Identification of essential oils components by gas chromatography/mass spectroscopy. Carol Stream, IL, USA: Allured Publication Corporation.
- [4]. Angioni, A.; Barra, A.; Arlorio, M.; Coisson, J.D.; Russo, M.T.; Pirisi, F.F., et al. (2003). Chemical composition, plant genetic differences, and antifungal activity of the essential oil of *Helichrysum italicum* G. Don ssp. *microphyllum* (Willd) Nym. *Journal of Agricultural and Food Chemistry*, 51, 1030-1034.
- [5]. Apel, K.; Hirt, H. (2004). Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annual Review of Plant Biology*, 55, 373-399.
- [6]. Ahmed M. (2012). Aboul-Enein, Faten Abu El-Ela, Emad A. Shalaby and Hany A. El-Shemy. Traditional medicinal plants research in Egypt: Studies of antioxidant and anticancer activities. *Journal of Medicinal Plants Research*, 6 (5), 689-703.
- [7]. Bakkali, F.; Averbeck, S.; Averbeck, D.; Idaomar, M. (2008). Biological effects of essential oils – A review. *Food and Chemical Toxicology*, 46, 446-475.
- [8]. Bora, K.S.; Sharma, A. (2011). The genus *Artemisia*: A comprehensive review. *Pharmaceutical Biology*, 49, 101-109.
- [9]. Brand-Williams, W., Cuvelier, M.E., Berset, C. (1995). Use of a free radical method to evaluate antioxidant activity. *LWT-Food Science and Technology*, 28, 25-30.
- [10]. Cai, Y.; Luo, Q.; Sun, M.; Corke, H. (2004). Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Science*, 74, 2157-2184.
- [11]. Daise Lopes-Luts, Daniela S.Alviano, Celuta S. Alviano, Paul P.Kolodziejczyk. (2008). Screening of chemical composition, antimicrobial and antioxidant activities of *Artemisia* essential oils. *Phytochemistry*, 69, 1732-1738.
- [12]. El-Shemy H.A.; Aboul-Enein, A.M.; Aboul-Enein, K.M.; Fujita, K. (2007). Willow leaves' extracts contain anti-Tumor agents effective against three cell types. *PLOS ONE*, 2(1), 178.
- [13]. Ferrigni, N.; Meyer, B.; Ferrigni, N.; Putnam, J.; Jacobsen, L.; Nichols, D.; Melaughlin, J. (1982). Brine shrimp: A convenient general bioassay for active plant constituents. *Journal of Planta Medica*, 45, 31-34.
- [14]. Grubov, V.I. (1982). Key to the Vascular Plants of Mongolia. Nauka: Leningrad, Russia. 245-253.
- [15]. Jian-Qing Yu, Jia-Chuan Lei, Xiu-Qiao Zhang, Huai-Dong Yu, Dai-Zhi Tian, Zhi-Xiong Liao, Guo-lin Zou. (2011). Anticancer, antioxidant and antimicrobial activities of the essential oil of *Lycopus lucidus* Turcz. var. *hirtus* Regel. *Food chemistry*, 126, 1593-1598.
- [16]. Kalemba, D., Kunicka, A. (2003). Antibacterial and antifungal properties of essential oils. *Current Medicinal Chemistry*, 10, 813-829.
- [17]. Markova, L.P. et al. (1985). Wild growing useful plants in flora Mongolia. Leningrad. 22, 89, 92.
- [18]. Lawrence, B.M., (2005). Antimicrobial/Biological activity of Essential Oils. Allured Publishing Corporation, Illinois, United States.
- [19]. Lee, J.; Godon, C.; Lagniel, G.; Spector, D.; Garin, J.; Labarre, J.; Toledano, M. B. (1999). Yap1 and Skn7 control two specialized oxidative stress response regulons in yeast. *Journal of Biological Chemistry*, 274 (23), 16040.
- [20]. Ligaa, U., Davaasuren, B., Ninjil, N. (2005). Using the Mongolian medicinal herbs in eastern and western medicine. KSA Press, UB, Mongolia, 114.
- [21]. Martinez-Tome, M., Jimenez, A., Ruggieri, S., Frega, N., Strabbioli, R., et al. (2001). Antioxidant properties of Mediterranean spices compared with common food additives. *Journal of Food Protection*, 64, 1412-1419.
- [22]. Massada. Y. (1976). Analysis of Essential Oil by Gas Chromatography and Spectrometry. Wiley, New York, USA.

- [23]. Molyneux, P. (2004). The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarin Journal of Science and Technology*, 26, 211-219.
- [24]. Moon, J.K.; Shibamoto, T. (2009). Antioxidant assays for plant and food components. *Journal of Agricultural and Food Chemistry*, 57(5), 1655-1666.
- [25]. Mosman, T. (1983). Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assay. *Journal of Immunological Methods*, 65, 55-63.
- [26]. Sylvestre, M., Pichette, A., Longtin, A., Nagau, F., & Legault, J. (2006). Essential oil
- [27]. analysis and anticancer activity of leaf essential oil of *Croton flavens* L. From
- [28]. Guadeloupe. *Journal of Ethnopharmacology*, 103, 99-102.
- [29]. Salzer, U.J. (1977). The analysis of essential oils and extracts (oleoresins) from seasonings – A critical review. *CRC Critical Reviews Food Science and Nutrition*, 9, 345-373.
- [30]. Senatore, F., Arnold, N.A., Piozzi, F. (2004). Chemical composition of the essential oil of *Salvia multicaulis* Vahl. var. *simplicifolia* Boiss. growing wild in Lebanon. *Journal of Chromatographia*, 1052, 237-240.
- [31]. Shatar, S. (1989). Mongolian fragrant herbs. Soyombo Press, UB, Mongolia. 14-16.
- [32]. Shatar. S., Altantsetseg, Sh. (2000). Essential oil composition of some plants cultivated in Mongolian climate. *Journal of Essential Oil Research*, 12, 745-750.
- [33]. Sikkema, J., J.A. de Bont, Poolman, B. (1995). Mechanisms of membrane toxicity of hydrocarbons. *Microbiological Reviews*, 59, 201-222.
- [34]. Sylvester, P.W. (2011). Optimization of the tetrazolium dye (MTT) colorimetric assay for cellular growth and viability. *Methods Molecular Biology*, 716, 157-68.
- [35]. Tang, S. Y.; Whiteman, M.; Peng, Z. F.; Jenner, A.; Yong, E. L.; Halliwell, B. (2004). Characterization of antioxidant and antiglycation properties and isolation of active ingredients from traditional Chinese medicines. *Free Radical Biology and Medicine*, 36, 1575-1587.
- [36]. Valko, M.; Leibfritz, D.; Moncol, J.; Cronin, M.; Mazur, M.; Telser, J. (2007). Free radicals and antioxidants in normal physiological functions and human disease. *International Journal of Biochemistry and Cell Biology*, 39, 44-84.
- [37]. Vandendool, H., Kratz, P.D. (1963). A generalization of the retention index system including linear temperature programmed gas liquid partition chromatography. *Journal of Chromatographia*, 11, 463–471.
- [38]. Verezovskaya, T.P. et al. (1991). Worm woods in Siberia. *Novosibirsk*. 48, 87-96.
- [39]. Zhang, X. (2004). "Traditional medicine: its importance and protection. In: Twarog S, Kapoor P (eds). *Protecting and Promoting Traditional Knowledge: Systems, National Experiences and International Dimensions*. Part 1. *The Role of Traditional Knowledge in Healthcare and Agriculture*". New York: United Nations, 3-6.